

Locations:

☐ Dublin, OH☐ Lancaster, OH

PATIENT INFORMATION

Patient Name

Date of Birth

Gender

☐ M☐ F

Patient Address / City / State / ZIP

Phone

DIAGNOSIS

PROVIDE ICD-10 CODE

☐ Postmenopausal osteoporosis - high fracture risk☐ Male osteoporosis - high fracture risk☐ Glucocorticoid-induced osteoporosis - high fracture risk☐ Bone loss due to androgen-deprivation therapy☐ Bone loss due to adjuvant aromatase-inhibitor therapy☐ Other

PRE-MEDICATIONS

OPTIONAL - CHECK ALL THAT APPLY

☐ None☐ Acetaminophen 1,000 mg PO☐ Diphenhydramine 25 mg PO☐ Cetirizine 10 mg PO☐ Other

BILDYOS ORDERS

DOSAGE & FREQUENCY

Last denosumab injection date - if applicable

☐ 60 mg subcutaneous injection every 6 months☐ New start☐ Continuation

Injection site: upper arm, upper thigh, or abdomen.

REQUIRED CLINICAL REVIEW

COMPLETE BEFORE ADMINISTRATION

☐ Pregnancy ruled out, if applicable☐ Pre-existing hypocalcemia corrected☐ Serum calcium reviewed☐ Routine oral examination completed before initiation☐ If eGFR <30 or dialysis: CKD-MBD evaluation completed☐ Calcium 1,000 mg/day and vitamin D at least 400 IU/day instructed

NOTES

ORDERING PROVIDER

Provider NPI

Phone

Fax

Provider Name

Provider Address

Prescriber Signature

Date