



# Dermatology Referral Form

**\*\*Please Attach Copy of Insurance Cards (Front & Back)\*\***

Last Name, First Name: _____	Date of Birth: _____
Address: _____	Gender: <input type="checkbox"/> M <input type="checkbox"/> F <input type="checkbox"/> Other
Phone: _____	City, State, Zip: _____
Referring Practice: _____	_____
Practice Address: _____	City, State, Zip: _____
Prescriber Name: _____	Prescriber NPI: _____
Nurse/Key Contact: _____	Phone: _____
Fax: _____	Email: _____

## Nursing & Lab Orders

**Nurse Orders:** Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.  
**Flush Orders:** NaCl 0.9% - 5-10mL flush pre and post infusion and as needed *Heparin* -  10units/mL ---OR---  100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line  
**Lab Orders:** \_\_\_\_\_ **Lab Date & Frequency:** \_\_\_\_\_

## Prescription Orders

<b>Anaphylaxis Kit:</b>	<input type="checkbox"/> Epinephrine 0.3mg IM as needed	<input type="checkbox"/> Solu-cortef 250mg-500mg IV infusion as needed	<input type="checkbox"/> Solu-Medrol 60mg - 125mg IV infusion as needed
(Check all that apply)	<input type="checkbox"/> Diphenhydramine _____mg IV infusion as needed	<input type="checkbox"/> NS Hydration 500 ml IV infusion over 30 minutes as needed	<input type="checkbox"/> Other
<b>Pre-Medications:</b>	<input type="checkbox"/> Acetaminophen _____mg PO _____minutes prior to infusion	<input type="checkbox"/> Solu-Medrol _____mg IV _____minutes prior to infusion	
(Check all that apply)	<input type="checkbox"/> Diphenhydramine _____mg <input type="checkbox"/> PO ---OR--- <input type="checkbox"/> IV infusion _____minutes prior to infusion	<input type="checkbox"/> Other	
<b>Supply Orders:</b> All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary			

## Prescription Information

PRODUCT	DIRECTIONS	REFILLS
Is this a first dose? <input type="checkbox"/> Yes <input type="checkbox"/> no if no, when was last dose given? _____ When is patient due for next dose? _____		
<input type="checkbox"/> Ilumya	100mg sc injection at 0 and 4 weeks then every 12 weeks	
<input type="checkbox"/> Infliximab <input type="checkbox"/> Avsola <input type="checkbox"/> Inflectra <input type="checkbox"/> Remicade <input type="checkbox"/> Renflexis	<input type="checkbox"/> <b>INDUCTION:</b> _____mg/kg or _____mg iv infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over at least 2 hours at weeks 0, 2, and 6	NONE
	<input type="checkbox"/> <b>MAINTENANCE:</b> _____mg/kg _____mg iv infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over at least 2 hours every _____ weeks (Note: round to nearest 100mg for medicaid patients) If remicade infusion tolerated, adjust infusion time according to manufacturer package insert.	
<input type="checkbox"/> Simponi Aria	2 mg/kg IV infusion via <input type="checkbox"/> gravity ---OR--- <input type="checkbox"/> pump over 30 minutes at weeks 0 and 4, and every 8 weeks thereafter	
<input type="checkbox"/> Spevigo	<input type="checkbox"/> 900 Mg iv infusion over 90 minutes <input type="checkbox"/> Additional 900 mg iv infusion over 90 minutes one week after initial dose if flare symptoms persist	
<input type="checkbox"/> Stelara	<b>Psoriasis Adult Subcutaneous</b> <input type="checkbox"/> For patients <= 100 kg, 45 mg SC injection initially and 4 weeks later, followed by 45 mg every 12 weeks <input type="checkbox"/> For patients > 100 kg, 90 mg SC injection initially and 4 weeks later, followed by 90 mg every 12 weeks	
	<b>Psoriasis Pediatric Patients 6 to 17 (based on weight at time of initial dose)</b> <input type="checkbox"/> For patients <= 60 kg, 0.75 mg/kg SC injection initially and 4 weeks later, then every 12 weeks <input type="checkbox"/> For patients 60 kg – 100kg, 45 mg SC injection initially and 4 weeks later, then every 12 weeks <input type="checkbox"/> For patients >100kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks	
	<b>Psoriasis Adult Subcutaneous</b> <input type="checkbox"/> 45 mg SC injection initially and 4 weeks later, followed by 45 mg SC injection every 12 weeks <input type="checkbox"/> For patients with co-existent moderate-to-severe plaque psoriasis weighing >100 kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks	
<input type="checkbox"/> Xolair	<input type="checkbox"/> 150 or <input type="checkbox"/> 300 mg SC injection once every 4 weeks	
<input type="checkbox"/> IG	<b>For Immunoglobulin therapy please refer to Immunoglobulin Form</b>	
<input type="checkbox"/> Other		

**By signing this form and utilizing our services, you are authorizing Infuse One Ohio to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.**

### DISPENSE AS WRITTEN

Prescriber's Signature: \_\_\_\_\_ Print Name: \_\_\_\_\_ Date: \_\_\_\_\_

### SUBSTITUTION PERMITTED

Prescriber's Signature: \_\_\_\_\_ Print Name: \_\_\_\_\_ Date: \_\_\_\_\_

**Please be sure to attach the following:** Provider Phone Line: (614) 929-3349

- |   |   |
|---|---|
| <input type="checkbox"/> Patient demographics & front/back copy of all insurance cards (prescription & medical) | <input type="checkbox"/> TB lab results within last 12 months ( <i>Stelara, Simponi Aria, Ilumya &amp; Infliximabs only</i> ) |
| <input type="checkbox"/> Recent office visit notes, history & physical, lab & pertinent procedure results       | <input type="checkbox"/> HBV lab results within last 12 months ( <i>Infliximabs &amp; Simponi Aria only</i> )                 |
| <input type="checkbox"/> Current medication list & list of prior medications tried and failed (with dates)      | <input type="checkbox"/> Letter of medical necessity if drug dosing or indication is outside of FDA guidelines                |